

WHY GALAXIES CARE ABOUT AGB STARS III:
A CLOSER LOOK IN SPACE AND TIME
ASP Conference Series, Vol. 497
Kerschbaum, Wing, and Hron, eds.
©2015 Astronomical Society of the Pacific

Where is the Metallicity Ceiling to Form Carbon Stars? – A Scarcity of Carbon Stars in the Inner Disk of M31

Martha L. Boyer,¹ L. Girardi,² P. Marigo,³ B. F. Williams,⁴ B. Aringer,⁵
W. Nowotny,⁵ P. Rosenfield,^{3,4} C. E. Dorman,⁶ P. Guhathakurta,⁶
J. J. Dalcanton,⁴ J. L. Melbourne,⁷ K. A. G. Olsen,⁸ and D. R. Weisz⁶

¹*Observational Cosmology Lab, NASA Goddard Space Flight Center,
Greenbelt MD, USA*

²*INAF – Osservatorio Astronomico di Padova, Italy*

³*Dept. of Physics and Astronomy G. Galilei, University of Padova, Italy*

⁴*Dept. of Astronomy, University of Washington, Seattle WA, USA*

⁵*Dept. of Astrophysics, University of Vienna, Austria*

⁶*University of California Observatories/Lick Observatory, Santa Cruz CA, USA*

⁷*Caltech Optical Observatories, Pasadena CA, USA*

⁸*National Optical Astronomy Observatories, Tucson AZ, USA*

Abstract. We have used a novel technique involving near-infrared colors from the Hubble Space Telescope to distinguish C stars from M stars in a field in M31. The resulting C/M ratio is much lower than other estimates in M31.

Carbon stars are far more efficient at producing dust than O-rich (M-type) Asymptotic Giant Branch (AGB) stars (e.g. Matsuura et al. 2009; Srinivasan et al. 2009), and may be the dominant stellar dust source in galaxies (e.g. Boyer et al. 2012; Zhukovska & Henning 2013). They are also responsible for a major share of galaxy luminosity (e.g. Melbourne & Boyer 2013) and contribute significantly to the chemical enrichment of the interstellar medium (e.g. Karakas & Lattanzio 2007; Marigo 2001; Ventura et al. 2001). Because of their vital role in galaxy evolution and in interpreting the observations of unresolved galaxies, it is important to devise a method to efficiently identify C stars in distant systems that will become resolvable with new large telescopes.

In Boyer et al. (2013), we use medium-band near-infrared (NIR) Hubble Space Telescope WFC3 photometry with model NIR spectra of Asymptotic Giant Branch (AGB) stars to develop a new tool for efficiently distinguishing carbon-rich (C-type) AGB stars from oxygen-rich (M-type) AGB stars in galaxies at the edge of and outside the Local Group (Figure 1). We test this technique in a $2'.3 \times 2'.1$ field of the inner disk of M31, 2 kpc from the center, where several C star candidates were detected via their *JHK* colors (Davidge et al. 2005). We detect only 1 candidate C star (plus up to 6 additional, less certain C stars candidates), resulting in an extremely low ratio of C to M stars ($C/M = 3.3^{+20}_{-0.1} \times 10^{-4}$) that is ~ 2 orders of magnitude lower than other C/M estimates in M31 (Brewer et al. 1995).




Figure 1. *Left:* The $F139M-F153M$ and $F127M-F139M$ color combination efficiently separates C and M type AGB stars with little to no overlap. In addition, M type stars follow a sequence of increasing H_2O absorption, and thus temperature. *Right:* The same for the observed region of M31. Only one candidate C star is detected with high confidence. The candidate C stars from Davidge et al. (2005) are instead late-type M stars with deep H_2O absorption. H_2O absorption causes red JHK colors; caution is required when using standard NIR filters to identify C stars.

The low C/M ratio is likely due to the high metallicity in this region which impedes stars from achieving $C/O > 1$ in their atmospheres (Marigo et al. 2013). Previous C star identifications in this region using JHK colors are likely misidentified M stars with deep water absorption near $1.4 \mu m$. These observations provide stringent constraints to evolutionary models of metal-rich AGB stars and point to a metallicity threshold above which M stars are unable to make the transition to C stars, dramatically affecting AGB mass loss and dust production and, consequently, the observed global properties of metal-rich galaxies. The metallicity of our field suggests that the ceiling for C stars falls in the range $-0.1 < [M/H] < +0.1$.

Acknowledgments. This work was supported by the NASA Postdoctoral Program at the GSFC, administered by ORAU through a contract with NASA, and by NASA through HST grant numbers GO-12862 and GO-12055 from the STScI, which is operated by AURA, Inc., under NASA contract NAS5-26555. PM and LG acknowledge support from Progetto di Ateneo 2012, University of Padova, ID: CPDA125588/12. This research was funded in part by the Austrian Science Fund (FWF): P21988-N16.

References

- Aringer, B., Girardi, L., Nowotny, W., Marigo, P., & Lederer, M. T. 2009, *A&A*, 503, 913
- Boyer, M. L., Girardi, L., Marigo, P., et al. 2013, *ApJ*, 774, 83
- Boyer, M. L., Srinivasan, S., Riebel, D., et al. 2012, *ApJ*, 748, 40
- Brewer, J. P., Richer, H. B., & Crabtree, D. R. 1995, *AJ*, 109, 2480
- Davidge, T. J., Olsen, K. A. G., Blum, R., Stephens, A. W., & Rigaut, F. 2005, *AJ*, 129, 201
- Karakas, A., & Lattanzio, J. C. 2007, *PASA*, 24, 103
- Marigo, P. 2001, *A&A*, 370, 194
- Marigo, P., Bressan, A., Nanni, A., Girardi, L., & Pumo, M. L. 2013, *MNRAS*, 434, 488
- Matsuura, M., Barlow, M. J., Zijlstra, A. A., et al. 2009, *MNRAS*, 396, 918
- Melbourne, J., & Boyer, M. L. 2013, *ApJ*, 764, 30
- Srinivasan, S., Meixner, M., Leitherer, C., et al. 2009, *AJ*, 137, 4810
- Ventura, P., D’Antona, F., Mazzitelli, I., & Gratton, R. 2001, *ApJL*, 550, L65
- Zhukovska, S., & Henning, T. 2013, *A&A*, 555, A99